

COMPARATIVE STUDY OF THE MUON-CATALYZED
FUSION IN D-T AND D-³He SYSTEMS

L. F. Hansen
University of California
Lawrence Livermore National Laboratory
Livermore, CA 94550

September 1985

Lawrence
Livermore
National
Laboratory

This is an informal report intended primarily for internal or limited external distribution. The opinions and conclusions stated are those of the author and may or may not be those of the Laboratory.
Work performed under the auspices of the U.S. Department of Energy by the Lawrence Livermore National Laboratory under Contract W-7405-Eng-48.

DISCLAIMER

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial products, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

Printed in the United States of America
Available from
National Technical Information Service
U.S. Department of Commerce
5285 Port Royal Road
Springfield, VA 22161
Price: Printed Copy \$ Microfiche \$4.50

<u>Page Range</u>	<u>Domestic Price</u>	<u>Page Range</u>	<u>Domestic Price</u>
001-025	\$ 7.00	326-350	\$ 26.50
026-050	8.50	351-375	28.00
051-075	10.00	376-400	29.50
076-100	11.50	401-426	31.00
101-125	13.00	427-450	32.50
126-150	14.50	451-475	34.00
151-175	16.00	476-500	35.50
176-200	17.50	501-525	37.00
201-225	19.00	526-550	38.50
226-250	20.50	551-575	40.00
251-275	22.00	576-600	41.50
276-300	23.50	601-up ¹	
301-325	25.00		

¹Add 1.50 for each additional 25 page increment, or portion thereof from 601 pages up.

TABLE OF CONTENTS

I. RESEARCH PROPOSAL	
a) Abstract	01
b) Introduction	03
c) Measurements	06
d) Analysis of the Results	08
e) Acknowledgements	09
f) References	10
g) Figure Captions	12
II. TEACHING ACTIVITIES	13
III. RATIONALE FOR THE CHOICE OF HOST INSTITUTION	14
IV. RELATION OF THIS PROJECT TO NSF-VPS PROGRAM	15

I RESEARCH PROPOSAL

COMPARATIVE STUDY OF THE MUON-CATALYZED FUSION IN D-T AND D-³He SYSTEMS

L. F. Hansen

Lawrence Livermore National Laboratory. Livermore, CA 94550

ABSTRACT

In the last five years there has been a revival of interest in muon (μ)-catalyzed fusion reactions as a source of energy production. The measurements and calculations reported¹ deal mainly with the D-D and D-T systems which result in an energy release of 3.5 and 17.6 MeV respectively. The secondary neutrons produced in these reactions carry the larger fraction of the energy, 2.3 MeV in the D-D and 14.1 MeV in the D-T reactions. The study of the muon-catalyzed fusion of the light-ion system D-³He is of interest since the absence of radioactive tritium makes this reaction a source of "Clean Energy Fuel". In this reaction, $d + {}^3\text{He} \rightarrow p + {}^4\text{He} + 18.3 \text{ MeV}$, a 14.7 MeV proton is emitted instead of the 14.1 neutron from the D-T system. Although these two mirror systems (isospin flip), D-T and D-³He, are equivalent in regard to the nuclear interaction (they differ only in the Coulomb interaction), it would be also of interest to study the differences among the mesomolecular processes that will give rise to the μ -catalyzed fusion.

The objective of this proposal is to carry out a comparative study of the muon-catalyzed reactions, D-T and D-³He by measuring the number of neutrons/ μ and protons/ μ under equivalent experimental conditions. The comparison will be made as function of the T or ³He concentration in the systems, their densities and temperatures. Since some measurements²⁻⁴ of the D-T system are already available, the present study will concentrate on the D-³He system and in that region of the parameter space (concentration-density-temperature) not covered by the existent neutron work, mainly, mixtures at very high pressures and temperatures.

The experimental results will be compared with calculations of the neutron^{5,6} and proton yields resulting from the mesomolecular processes taking place between the muon and the deuterium-tritium and deuterium-helium mixtures.

INTRODUCTION

The "Cold Fusion" process in which a catalyzing agent, such as a negative meson, induces the fusion reaction was first discussed by Frank⁷ (1947) and Sakharov⁸ (1948) and experimentally observed⁹ in 1957. The interest in this process as a potential energy source has been revived by recent experiments^{3,4}. Close to 100 fusions reactions have been measured for for each meson present in deuterium tritium mixtures at high density (\sim liquid hydrogen density, LHD). Independent of the viability of cold fusion as an economical choice for fusion power^{10,11}, the muon catalyzed fusion process is of interest because of the atomic, molecular and nuclear processes that are involved in its interpretation.

A consistent body of experimental results and further theoretical work are needed for an understanding of the muon-catalyzed fusion process with respect to the mesomolecular formation^{5,6} of the μDT , $\mu\text{D}^3\text{He}$ molecules. How does the strong nuclear interaction in the D-T, D- ^3He reactions affect¹² the molecular and atomic processes? How do the hyperfine components of the mesomolecular formation determine¹³ the fusion rates? How does the resonant theory of molecular formation hold at high temperatures? To answer these questions the appropriate measurements need to be performed. μ catalysis has been shown^{3,4} to depend on the density, temperature, and tritium concentration of the D-T mixture. This dependence needs to be systematically explored at higher pressures and temperatures. It is also of interest to extend this study to the D ^3He mixtures, for which calculations on the mesomolecular formation have only recently become available¹⁴.

The scope of the present proposal is to carry out measurements of the catalyzed fusion reaction for the D-T and the D- ^3He systems by observing the emission of neutrons and protons, respectively. These measurements, in collaboration with K. Crowe (UC, Berkeley), and C. Petitjean (Swiss Institute for Nuclear Research, SIN, Switzerland) will look first into the neutron emission as a function of temperature and pressure of the D-T mixture. Some

overlap with early experiments is desirable in order to test the reproducibility of the earlier results^{3,4}. However, the interest of the proposed measurements is principally into the regime of high pressures (~ 1000 atm) and temperatures ($1-5 \times 10^3$ K). Both Jones et al.³ and Breunlich et al.⁴ have shown an increasing, but different temperature dependence of the neutron production rate (per muon stopped in the D-T mixture) in the 30 to 900 K interval. The interpretation of the results differ among these groups. Observed⁴ hyperfine components of the μ td formation rate seem to require a revision of the predicted¹⁵ neutron rates and its dependence on temperature.

Measurements at higher temperatures ($1-5 \times 10^3$ K) would test the resonance theory of the μ td molecule formation. Would the system be "thrown out of resonance" at these temperatures and the catalysis process be stopped? Would there be a continuous increase of the neutron rates as a result of new resonances being excited? Similar questions could be answered by measurements at high pressures. Does the neutron rate increase progressively with higher densities of the D-T mixture or is a plateau reached at a given pressure? Does the catalysis stop at very high densities due to an increasing number of α particles and increasing μ ⁴He sticking?

Proton measurements will follow after replacing the D-T mixture by D-³He gas for the same experimental conditions of ³He concentration, pressure and temperature. The comparison of the rates of neutron and proton emission per incident μ meson from the catalyzed D-T and D-³He reactions would give interesting information into the mesomolecular processes between these two reactions. Would protons be observed in coincidence with the meson? In other words, is there a bound state for the μ ³HeD molecular system as in the D-T mixture? How different are the lifetimes of the μ t and μ ³He atoms? Would the sticking coefficient of the μ to the ³He be of the same order as the value obtained¹⁶ for the α particle? Are the mesomolecular processes affected by the isospin flip ($n \rightarrow p$) term in the strong interaction Hamiltonian for the $D + T$ and $D + {}^3\text{He}$ nuclear reactions?

The proposed measurements would shed light on the possibility of using the $D-^3\text{He}$ reaction as a source of catalyzed fusion. This is in itself a very attractive choice since it eliminates the need for tritium with its intrinsic health hazard problems. Furthermore, the comparative study of the neutron/proton emission would stimulate further theoretical work into the μTD system and needed calculations for the $\mu^3\text{HeD}$. Some insight also could be gained on the effect of nuclear processes from two equivalent (only isospin flip) nuclear reactions on the catalyzed fusion process.

MEASUREMENTS

The measurements will be done at SIN using the μ E4 beam and the experimental setup would be essentially the same used in previous measurements^{4,13}. A schematic view of the double wall target cell (a safety requirement for tritium handling) and detector system is shown in Fig. 1. For the high pressure D-T measurements (10^3 at), a new cell will be designed. It will be built at LLNL using their expertise in high pressure targets and tritium handling. For the D- ^3He mixture a simplified design (no need for a double wall container) would be used. Details of the gas pumping, storage and purification systems have been given elsewhere¹³.

A beam telescope consisting of three plastic scintillators, M_1 , M_2 , M_3 , defines the muon stops in the cell while the electrons from muon decay are detected by plastic scintillators E_1 , E_2 and E_3 (see Fig. 1). This arrangement^{4,13} of detectors and their electronics would be common to both the neutron and proton measurements.

Neutron Measurements

The nominal 14 MeV neutrons will be detected using standard¹⁷ time-of-flight (TOF) techniques. The detectors, NE213 liquid scintillators (30cm diam x 20cm) will be positioned one at each side of the gas cell. They will be viewed with 13 cm diam 8854RCA photomultipliers optically coupled to the detectors through a light pipe. Pulse-shape discrimination will be used to suppress the background. Neutron background from interactions of the muons or the 14 MeV fusion neutrons with the walls of the cell will be identified by TOF. The detector efficiency for 14 MeV neutrons will be obtained by a Monte Carlo calculation using the code¹⁸ NEFF4. Good agreement has been found¹⁹ with this code for the measured and calculated efficiencies for NE213 and Stilbene detectors. Multiple scattering corrections will need to be made to account for the thick walls of the cell and high gas pressure.

Proton Measurements

The proton measurements will be made first for the same range of pressures ($1-3 \times 10^3$ atm) and temperatures (10-900 K) as the neutron measurements (See summary of the existing measurements in Pg. 55 of Ref. 1). Only at very low pressures (a few atmospheres) would it be possible to have a thin window, so that the energy lost by the 14.6 MeV proton in the mixture and through the window will not be larger than a few hundred keV. This would allow the proton detector to be outside the gas cell. Silicon surface barrier detectors ($\Delta E-E$ telescope), 150-200 microns total thickness, would be used to detect the protons. Because of the smaller diameter of the Si detectors, the solid angle subtended by the proton telescope could be about an order of magnitude smaller than that of the neutron detector if installed at the same position. However, by bringing the detector close to the window and because of the larger efficiency of the Si detectors (100% versus 3-4% for the neutron detector) an overall gain in detection efficiency is obtained.

For measurements at very high pressures the Si detectors would be installed inside the gas cell (passivated Si detectors with an epitaxial layer over the p-n junction) because of the large fraction of energy lost by the proton in the $D-^3He$ mixture. i.e. for a mixture at 1000 atm pressure the total energy lost by the 14.6 MeV proton in reaching the walls of a 5cm diameter cell is around 9 MeV. Furthermore, at these high pressures the thickness of the window will stop altogether the already low-energy emerging protons.

To install the proton detectors inside the gas cell is a very practical approach that could be used over the entire range of pressures of the measurements. The only limitation would be as a result of the increasing leakage current of the Si detectors at higher temperatures. Under this arrangement, 450-500 K could be an upper limit for these measurements.

ANALYSIS OF THE RESULTS

The analysis of the data will be carried out at Berkeley. The Monte Carlo calculations of the neutron detector efficiency and of the neutron and proton transmission from the gas cell to the detectors will be done using the 7600 MFI computer. The nuclear cross sections for the $T(d,n)^4\text{He}$ and $^3\text{He}(d,p)^4\text{He}$ reactions in the eV region and their angular distributions would be obtained from an R-matrix analysis²⁰ of the mass-5 system.

The measured neutron emission rates, after being corrected for background contributions from neutrons originating in the cell walls, will be compared with calculations. The results will be compared with the predicted decay rates of the mesomolecular formation as obtained from previous analyses^{4,13}, in order to clarify the role of the hyperfine structure decay channels. Fig. 2 is illustrative of the model for the muon catalysis in the DT mixture as presently accepted¹³.

A parallel analysis of the proton emission rates would be a test of the recent Russian calculations¹⁴ and would give information as to the applicability of the model to the $D^3\text{He}$ system. Calculations have been initiated^{21,22} based on this proposal on the mesomolecular formation mechanism for the $\mu^3\text{HeD}$ molecule and the values for the sticking coefficients of the μ to the ^3He and ^4He atoms. It is expected that these calculations will be available at the time of completion of these measurements.

Note added in proof:

Although the term "mesomolecular" has been used throughout the text, according to its common usage in the literature, it is a poor choice²³ since the muon is not a meson.

ACKNOWLEDGEMENTS

The author would like to acknowledge stimulating discussions with K. Crowe, D. Ceperly and M. S. Weiss.

REFERENCES



1. "Muon- Catalyzed Fusion Workshop", June 7-8, 1984. Jackson Hole, Wyoming. Steven E. Jones, Editor. EG&G, Idaho Falls, Idaho 83415. See papers presented to this workshop and their references.
2. V. M. Bystritsky et al., Phys. Lett. 94B, 476 (1980).
3. Steven E. Jones, et al., Atomkernenergie-Kerntechnik, 43 (3), 179 (1983); Phys. Rev. Lett. 51, 1757 (1983).
4. W. H. Breunlich et al., Phys. Rev. Lett., 53, 1137 (1984).
5. S. I. Vinitiskii, et al., Sov. Phys. JETP 47 (3), 444 (1978)
6. L. N. Bogdanova, et al., Sov. Phys. JETP 34 (5), 662 (1981); 54 (3), 442 (1981); 56 (5), 931 (1982).
7. F. C. Frank. Nature 160, 525 (1947).
8. A. D. Sakharov, Report of the Phys. Inst., Academy of Sci., USSR, (1948).
9. L. W. Alvarez et al., Phys. Rev. 105, 1127 (1957).
10. T. K. Larson et al., " Analysis of System Requirements for a Commercially Feasible Muon-Catalyzed Fusion Power Plant". 3rd Int. Conf. on Emerging Nucl. Systems. Helsinki, Finland, June 6-9 (1983).
11. V. Yu. Petrov, Nature 285, 466 (1980).
12. V. B. Belyaev, et al., J. Phys. G: Nucl. Phys. 8, 903 (1982).

13. P. Kammel et al., Atomkernenergie-Kerntechnik, 43, 195 (1983). Phys. Rev. A 28, 2611 (1983).
14. A. V. Kravtsov, A. I. Mikhailov, N. P. Popov, LNPI, Leningrad, Preprint-998, September 1984. See also Prep.-958, June 1984.
15. A. V. Matveenko et al., Sov. Phys. JETP 32, 871 (1971).
16. D. Ceperly and B. J. Alder, Phys. Rev. A 31, 1999 (1985).
17. L. F. Hansen et al., Phys. Rev. C 31, 111 (1985).
18. G. Dietze, private communication February (1985). Phys.-Tech. Bundesanstalt, Braunschweig.
19. L. F. Hansen et al., UCRL-92426, April 1985. Accepted for publication in Nucl. Sci. & Eng. See Fig. 2 and other refs.
20. G. M. Hale, Los Alamos National Laboratory, Private Communication, May (1985).
21. D. Ceperly, Lawrence Livermore National Laboratory, Private Communication, September (1985).
22. M. S. Weiss and Chi-Yu Hu, Lawrence Livermore National Laboratory Private Communication, September (1985).
23. L. Bracci and G. Fiorentini, Phys. Reports 86, 169 (1982).

FIGURE CAPTIONS

Fig. 1 Schematic representation of the experimental arrangements for the D-T (Fig. 1a) and D-³He (Fig. 1b) mixtures. M₁, M₂, M₃ muon beam defining detectors. E₁, E₂, E₃ muon's decay electron detectors. NE213 neutron detectors. P₁, P₂ proton detectors

Fig. 2 Model of the processes taken place when negative muons are stopped in a mixture of deuterium and tritium, including hyperfine transitions. (Taken from Ref. 4). c_d and c_t are the atomic concentrations of deuterium and tritium. The λ's correspond to the different rates of molecular formation. λ_{hf} denotes the hyperfine transition: $\lambda_{hf} = c_t \lambda_{tt}^{\mu} + c_d \lambda_{dt}^{\mu}$. J is the total spin of the μt atom and w_s is the sticking coefficient of the muon to the ⁴He.






Fig. 2. The DT fusion cycle